

To appear in *Nature*, 8 January 1998.

Further evidence for the planet around 51 Pegasi

Artie P. Hatzes, William D. Cochran, & Eric J. Bakker

McDonald Observatory, The University of Texas at Austin, Austin, TX 78712

The discovery¹ of the planet around the solar-type star 51 Pegasi marked a watershed in the search for extrasolar planets. Since then seven other solar-type stars have been discovered^{2–6}, of which several have surprisingly short orbital periods, like the planet around 51 Peg. These planets were detected using the indirect technique of measuring variations in the Doppler shifts of lines in the spectra of the primary stars. But it is possible that oscillations of the stars themselves (or other effects) could mimic the signature of the planets, particularly around the short-period planets. The apparent lack of spectral⁷ and brightness variations⁸, however, led to widespread acceptance that there is a planet around 51 Peg. This conclusion was challenged by the observation⁹ of systematic variations in the line profile shapes of 51 Peg, which suggested stellar oscillations¹⁰. If these observations are correct, then there is no need to invoke a planet around 51 Peg to explain the data. Here we report observations of 51 Peg at a much higher spectral resolution than those in ref. 9, in which we find no evidence for systematic changes in the line shapes. The data are most consistent with a planetary companion to 51 Peg.

Observations were made using the coudé echelle spectrograph¹¹ of the 2.7-m Harlan J. Smith telescope at the McDonald Observatory. This instrument was used in the configuration that provided a resolving power ($\lambda/\Delta\lambda$) of 220,000, or more than twice that of the data obtained by Gray⁹, and with a wavelength coverage of several hundred (non-contiguous) Ångströms. Approximately 120 observations were made over the course of 18 nights in July – September 1997. On each night 4–10 observations of 51 Peg were made with a typical signal-to-noise ratio of ~ 200 for each spectrum. Information about the spectral line shapes was extracted using the line bisector which is the locus of the midpoints of the stellar absorption line from the line core up to the continuum. The velocity span and the bisector curvature were both measured. The span is the velocity difference between the bisector points taken at flux levels of 0.48 and 0.85 of the continuum level and the curvature is the difference in the velocity span between the top half (measured between flux points 0.64 and 0.85 of the continuum) and bottom half (measured between flux points 0.42 and 0.64 of the continuum) of the line bisector. (See Gray⁹ and Gray & Hatzes¹⁰ for illustrations of a typical bisector as well as definitions of the velocity span and curvature.) Both the

span and the bisector curvature were examined because it is possible for spectral variations to have large changes in the curvature without significant changes in the velocity span and vice-versa. The spectral coverage of our data included the 6,252.57-Å Fe I line used by Gray as well as 4 other spectral features of comparable line strength that were suitable for bisector measurements: 5,296.7-Å Cr I, 5,554.93-Å Fe I, 5,639.95-Å Fe I, and 6141.75-Å Fe I. If the kinematic nonradial pulsation model proposed by Gray & Hatzes¹⁰ is correct, then changes in the lines bisectors for spectral lines of equal strength should be the same, assuming that the spectral lines are formed at the same depth in the stellar atmosphere. Averaging bisectors from several line should decrease the noise level in the measurements.

Figure 1 shows the results of our velocity span measurements. The top panel shows the nightly average of the velocity span of 6,252.57-Å Fe I, the same feature measured by Gray⁹, as a function of phase. Phases were computed using the orbital ephemeris of Marcy et al.¹² (a period of 4.2311 d and an epoch of Julian day 2449797.773) which was also used by Gray and Hatzes¹⁰. The solid line represents the predicted sinusoidal variation of Gray⁹. Each error bar is the standard deviation of individual nightly measurements weighted by the square root of the number of measurements used in the average. The r.m.s. scatter of the measurements about the mean is $\sim 25 \text{ m s}^{-1}$, which is half of the amplitude of the variations found in 51 Peg by Gray⁹ and is comparable to the r.m.s. scatter for non-variable stars. There may be a hint of sinusoidal variability in the measurements, but 180° out-of-phase with the curve predicted by Gray which suggests that this variability is due to noise. This is confirmed in the lower panel which shows the average velocity span for all five spectral lines that were examined. The errors shown represent the weighted standard deviation of the span measurements from the five spectral lines. The scatter (standard deviation) of the points shown is $\sim 7 \text{ m s}^{-1}$, well below the amplitude of the Gray prediction (line).

Figure 2 shows the result of our bisector curvature measurements. Once again, the top panel is for the 6,252.57-Å Fe I feature and the bottom panel is the average for all spectral lines. Errors were computed in the same manner as for Figure 1. The curve represents the fit to the variations from Gray & Hatzes¹⁰. Spectral variations, above the noise, do not seem to be present in either the measurements for the individual 6,252.57-Å Fe I or in the average curvature for all 5 spectral lines. The standard deviation of the averaged measurements (lower panel) is about 12 m s^{-1} .

The best direct comparison of our data to Gray’s is made with the 6,252.57-Å Fe I, the only featured in common with both data sets. Unfortunately, these alone cannot refute the Gray result with complete certainty. Phasing the data to the 4.2311 day radial velocity period does show a 180° phase shift between the two measurements and tentatively this would seem to contradict his claim of spectral variability. However, the scatter of our

measurements are comparable to the amplitude of the variations which may be present. A periodogram analysis using our data combined with Gray's shows this period as the highest peak, and the inclusion of our data reduces the power at secondary peaks. We note, however, that a periodogram analysis (not shown) of our data only does not show any significant power (false alarm probability $\sim 50\%$) at the appropriate period. A more detailed analysis and its implications will be presented elsewhere.

The averaged bisector measurements, on the other hand, almost certainly refute the Gray result, but not without caveats. Using the averaged measurements from different lines is valid so long as the spectral variations are the same (and in phase) for all spectral lines used in the analysis. This is the case for the kinematic pulsation model examined by Gray & Hatzes.¹⁰ which is only true if all spectral lines are formed at the same depth of the stellar atmosphere and have similar excitation potentials. However, only the 5,296.70Å Cr I line has a significantly different excitation potential (0.98eV) than the other spectral lines (3.6-4.5eV). If 51 Pegasi is really a long period pulsating star, then the vertical (radial) wavelength of the pulsations should be very short and spectral lines formed at different depths of the stellar atmosphere may show different behavior with phase in the spectral line shapes. If our line averaging technique is valid, then the spectral variability reported by Gray is most likely an artifact of noise. (Indeed, a Fourier analysis of the data in ref. 9 showed that there was about a 1 in 300 chance that pure noise could produce his observed variability.¹⁰) As the spectral variability of 51 Peg is not confirmed, the nonradial pulsation model for this star proposed by Gray & Hatzes¹⁰ is wrong. We will present elsewhere a more detailed analysis of our work, including line depth ratio measurements, variability of which was also reported by Gray⁹.

Stellar RV measurements provide only an indirect means of finding extra-solar planets and until there is a direct detection (either through imaging or from the spectral signature of the planet) we can never be absolutely certain that 51 Peg indeed has a planet. But in the light of all observational evidence now available for 51 Peg, including the measurements presented here, the planet hypothesis is the simplest and most plausible explanation for the variability reported by Mayor & Queloz¹.

Received 13 November, accepted 25 November 1997.

REFERENCES

1. Mayor, M. & Queloz, D. A Jupiter-mass companion to a solar-type star. *Nature*, **378**, 355-359 (1995).
2. Marcy, G.W. & Butler, R.P. A planetary companion to 70 Virginis *Astrophys. J.* **464**, L147-L151 (1996).
3. Butler, R.P. & Marcy, G.W. A planet orbiting 47 Ursae Majoris. *Astrophys. J.* **464**,

- L153-L156 (1996).
4. Butler, R.P., Marcy, G.W., Williams, E. Hauser, H. & Shirts, P. Three new ‘51 Peg-type’ planets. *Astrophys. J.* **474**, L115-L (1997).
 5. Cochran, W.D., Hatzes, A.P., Butler, R.P., & Marcy, G.W. The discovery of a planetary companion to 16 Cygni B. *Astrophys. J.* **483**, 457- (1997).
 6. Noyes, R.W., Jha, S., Korzennik, S.G., Krockenberger, M., Nisenson, P., Brown, T.M., Kennelly, E.J., Horner, S.D. A planet orbiting the star ρ Coronae Borealis. *Astrophys. J.* **483**, L111- (1997).
 7. Hatzes, A.P., Cochran, W.D., & Johns-Krull, C.M. Testing the planet hypothesis: a search for variability in the spectral line shapes of 51 Peg. *Astrophys. J.* **478**, 374- (1997).
 8. Henry, G.W., Baliunas, S.L. Donahue, R.A., Soon, W.H., Saar, S.H. *Astrophys. J.* **474**, 503- , (1997).
 9. Gray, D.F. Absence of a planetary signature in the spectra of the star 51 Pegasi. *Nature*, **385**, 795-796 (1997).
 10. Gray, D.F. & Hatzes, A.P. Nonradial oscillation in the solar-temperature star 51 Pegasi. *Astrophys. J.* **490**, 412–424 (1997).
 11. Tull, R.G., MacQueen, P.J., Sneden, C., & Lambert, D.L. The high-resolution cross-dispersed echelle white pupil spectrometer of the McDonald Observatory 2.7-m telescope. *PASP*, **107**, 251- (1995).
 12. Marcy, G.W., Butler, R.P., Williams, E., Bildsten, L., Graham, J.R., Ghez, A.M., Jernigan, J.G. The planet around 51 Pegasi. *Astrophys. J.* **481**, 926- (1997).




Fig. 1.— Velocity span measurements of 51 Peg. Top panel, the velocity span of 6,252.57 Fe I as a function of phase (data points) according to the ephemeris of Marcy et al.¹². The solid line represents the variations reported by Gray⁹. Bottom panel, the average velocity span for the spectral lines: 5,296.7-Å Cr I, 5,554.93-Å Fe I, 5,639.95-Å Fe I, 6141.75-Å Fe I, and 6,252.57 Fe I. Again, the line represents the predicted curve from Gray⁹.




Fig. 2.— Bisector curve measurements of 51 Peg. Top panel, the bisector curvature of 6,252.57 Fe I as a function of phase. The solid line represents the variations reported by Gray & Hatzes¹⁰. Bottom panel, the average bisector curvature for five spectral lines along with the predicted curve of Gray & Hatzes¹⁰.